

Symmetries in finite order variational sequences

Mauro Francaviglia, Marcella Palese¹, Turin,
Raffaele Vitolo², Lecce.

Abstract

We refer to Krupka's variational sequence, i.e. the quotient of the de Rham sequence on a finite order jet space with respect to a 'variationally trivial' subsequence. Among the morphisms of the variational sequence there are the Euler–Lagrange operator and the Helmholtz operator.

In this note we show that the Lie derivative operator passes to the quotient in the variational sequence. Then we define the *variational Lie derivative* as an operator on the sheaves of the variational sequence. Explicit representations of this operator give us some abstract versions of Noether's theorems, which can be interpreted in terms of conserved currents for Lagrangians and Euler–Lagrange morphisms.

Key words: Fibered manifold, jet space, variational sequence, symmetries, conservation laws, Euler–Lagrange morphism, Helmholtz morphism.

1991 MSC: 58A12, 58A20, 58E30, 58G05.

1 Introduction

The geometrical formulations of the Calculus of Variations on fibered manifolds include a large class of theories for which the Euler–Lagrange operator is a morphism of an exact sequence [1, 11, 17, 20, 21, 22]. This viewpoint allows one to overcome several problems of the Lagrangian formulations in Mechanics and Field Theories [19].

We consider the recent formulation by Krupka [11]. This has two main features. First, it is stated on finite order jets of the fibering, rather than on infinite order jets like most of the others. Moreover, it is conceptually very simple, because the exact sequence, or *variational sequence*, is defined as a quotient of the de Rham sequence on a finite order jet space with respect to an intrinsically defined subsequence, the choice of which is inspired by the Calculus of Variations.

¹ Partially supported by GNFM of CNR, MURST and University of Turin.

² Partially supported by GNFM of CNR, MURST, University of Florence and University of Lecce.

We show that the Lie derivative operator with respect to fiber-preserving vector fields passes to the quotient, yielding a new operator on the sheaves of the variational sequence, namely the *variational Lie derivative*. This idea has already been exploited by Vinogradov although in a different formalism [21, 22].

We make use of the representation given in [24] of the quotient sheaves of the variational sequence as concrete sheaves of forms. In this way, we provide explicit formulae for the quotient Lie derivative operators, as well as some abstract versions of Noether's theorems. Finally, we interpret Noether's theorems in terms of conserved currents for Lagrangians and Euler–Lagrange morphisms. In particular we recast, in a coherent and comprehensive scheme, some results previously found by various authors in different frameworks [1, 9, 10, 12, 13, 15, 18, 19].

We stress that the algebraic methods used here allow a synthetic and clear understanding of concepts whose full meaning could hardly be reached by means of coordinate expressions alone.

Throughout the paper the structure forms on jet spaces [14] as well as the horizontal and the vertical differential [16] will be used as fundamental tools.

Manifolds and maps between manifolds are C^∞ . All morphisms of fibered manifolds (and hence bundles) will be morphisms over the identity of the base manifold, unless otherwise specified. As for sheaves, we will use the definitions and the main results given in [26]. In particular, we will be concerned only with sheaves of \mathbb{R} -vector spaces. Thus, by ‘sheaf morphism’ we will shortly mean a morphism of sheaves of \mathbb{R} -vector spaces. Section 2 reviews earlier results, while Sections 3 and 4 contain the new results. An example is provided in Section 5.

Acknowledgments. Thanks are due to L. Fatibene, M. Ferraris, I. Kolář, D. Krupka and M. Modugno for useful comments. This work has been performed in the framework of Nat. Res. Proj. MURST 40% “Met. Geom. e Prob. in Fisica Matematica” and under the auspices of CNR–GNFM.

2 Jet spaces and variational sequences

In this section we recall some basic facts about jet spaces [3, 14, 16] and Krupka's formulation of the finite order variational sequence [11, 24].

2.1 Jet spaces

Here we introduce jet spaces of a fibered manifold and the sheaves of forms on the r -th order jet space. Moreover, we recall the notion of the horizontal and the vertical differential [16].

Our framework is a fibered manifold $\pi : \mathbf{Y} \rightarrow \mathbf{X}$, with $\dim \mathbf{X} = n$ and $\dim \mathbf{Y} = n + m$.

For $r \geq 0$ we are concerned with the r -jet space $J_r \mathbf{Y}$; in particular, we set $J_0 \mathbf{Y} \equiv \mathbf{Y}$. We recall the natural fiberings $\pi_s^r : J_r \mathbf{Y} \rightarrow J_s \mathbf{Y}$, $r \geq s$, $\pi^r : J_r \mathbf{Y} \rightarrow \mathbf{X}$, and, among

these, the *affine* fiberings π_{r-1}^r . We denote by $V\mathbf{Y}$ the vector subbundle of the tangent bundle $T\mathbf{Y}$ of vectors on \mathbf{Y} which are vertical with respect to the fibering π .

Charts on \mathbf{Y} adapted to π are denoted by (x^λ, y^i) . Greek indices λ, μ, \dots run from 1 to n and they label base coordinates, while Latin indices i, j, \dots run from 1 to m and label fibre coordinates, unless otherwise specified. We denote by $(\partial_\lambda, \partial_i)$ and (d^λ, d^i) the local bases of vector fields and 1-forms on \mathbf{Y} induced by an adapted chart, respectively.

We denote multi-indices of dimension n by boldface Greek letters such as $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_n)$ with $0 \leq \alpha_\mu$, $\mu = 1, \dots, n$; by an abuse of notation, we denote by λ the multi-index such that $\alpha_\mu = 0$ if $\mu \neq \lambda$, $\alpha_\mu = 1$ if $\mu = \lambda$. We also set $|\boldsymbol{\alpha}| := \alpha_1 + \dots + \alpha_n$ and $\boldsymbol{\alpha}! := \alpha_1! \dots \alpha_n!$.

The charts induced on $J_r\mathbf{Y}$ are denoted by (x^λ, y_α^i) , with $0 \leq |\boldsymbol{\alpha}| \leq r$; in particular, we set $y_0^i \equiv y^i$. The local vector fields and forms of $J_r\mathbf{Y}$ induced by the above coordinates are denoted by (∂_i^α) and (d_α^i) , respectively.

In the theory of variational sequences a fundamental role is played by the *contact maps* on jet spaces (see [3, 11, 12, 14]). Namely, for $r \geq 1$, we consider the natural complementary fibered morphisms over $J_r\mathbf{Y} \rightarrow J_{r-1}\mathbf{Y}$

$$\Pi : J_r\mathbf{Y} \times_{\mathbf{X}} T\mathbf{X} \rightarrow TJ_{r-1}\mathbf{Y}, \quad \vartheta : J_r\mathbf{Y} \times_{J_{r-1}\mathbf{Y}} TJ_{r-1}\mathbf{Y} \rightarrow VJ_{r-1}\mathbf{Y},$$

with coordinate expressions, for $0 \leq |\boldsymbol{\alpha}| \leq r-1$, given by

$$\Pi = d^\lambda \otimes \Pi_\lambda = d^\lambda \otimes (\partial_\lambda + y_{\alpha+\lambda}^j \partial_j^\alpha), \quad \vartheta = \vartheta_\alpha^j \otimes \partial_j^\alpha = (d_\alpha^j - y_{\alpha+\lambda}^j d^\lambda) \otimes \partial_j^\alpha.$$

We have

$$(1) \quad J_r\mathbf{Y} \times_{J_{r-1}\mathbf{Y}} T^*J_{r-1}\mathbf{Y} = \left(J_r\mathbf{Y} \times_{J_{r-1}\mathbf{Y}} T^*\mathbf{X} \right) \oplus \text{im } \vartheta_r^*,$$

where $\vartheta_r^* : J_r\mathbf{Y} \times_{J_{r-1}\mathbf{Y}} V^*J_{r-1}\mathbf{Y} \rightarrow J_r\mathbf{Y} \times_{J_{r-1}\mathbf{Y}} T^*J_{r-1}\mathbf{Y}$.

If $f : J_r\mathbf{Y} \rightarrow \mathbb{R}$ is a function, then we set $D_\lambda f := (\Pi)_\lambda f$, $D_{\alpha+\lambda} f := D_\lambda D_\alpha f$, where D_λ is the standard *formal derivative*. Given a vector field $Z : J_r\mathbf{Y} \rightarrow TJ_r\mathbf{Y}$, the splitting (1) yields $Z \circ \pi_r^{r+1} = Z_H + Z_V$ where, if $Z = Z^\gamma \partial_\gamma + Z_\alpha^i \partial_i^\alpha$, then we have $Z_H = Z^\gamma D_\gamma$ and $Z_V = (Z_\alpha^i - y_{\alpha+\gamma}^i Z^\gamma) \partial_i^\alpha$.

The splitting (1) induces also a decomposition of the exterior differential on \mathbf{Y} , $(\pi_r^{r+1})^* \circ d = d_H + d_V$, where d_H and d_V are defined to be the *horizontal* and the *vertical differential*. The action of d_H and d_V on functions and 1-forms on $J_r\mathbf{Y}$ uniquely characterizes d_H and d_V (see, e.g., [24] for more details).

A *projectable vector field* on \mathbf{Y} is defined to be a pair $(\Xi, \bar{\Xi})$, where $\Xi : \mathbf{Y} \rightarrow T\mathbf{Y}$ and $\bar{\Xi} : \mathbf{X} \rightarrow T\mathbf{X}$ are vector fields and Ξ is a fibered morphism over $\bar{\Xi}$. The coordinate expression of a projectable vector field is then $\Xi = \Xi^\lambda \partial_\lambda + \Xi^i \partial_i$, $\bar{\Xi} = \Xi^\lambda \partial_\lambda$, where Ξ^λ will depend only on coordinates on \mathbf{X} . If there is no danger of confusion, we will denote a projectable vector field $(\Xi, \bar{\Xi})$ simply by Ξ .

A projectable vector field $(\Xi, \bar{\Xi})$ can be conveniently prolonged to a projectable vector field $(j_r \Xi, \bar{\Xi})$, whose coordinate expression turns out to be

$$j_r \Xi = \Xi^\lambda \partial_\lambda + \left(D_\alpha \Xi^i - \sum_{\beta+\gamma=\alpha} \frac{\boldsymbol{\alpha}!}{\boldsymbol{\beta}! \boldsymbol{\gamma}!} D_\beta \Xi^\mu y_{\gamma+\mu}^i \right) \partial_i^\alpha,$$

where $\beta \neq 0$ and $0 \leq |\alpha| \leq r$ (see [10, 14, 16]); in particular, we have the expressions $(j_r \Xi)_H = \Xi^\lambda D_\lambda$, $(j_r \Xi)_V = D_\alpha (\Xi_V)^i \partial_i^\alpha$ with $(\Xi_V)^i = \Xi^i - y_\lambda^i \Xi^\lambda$. From now on, by an abuse of notation, we will drop the parentheses in $(j_r \Xi)_H$ and $(j_r \Xi)_V$ and write simply $j_r \Xi_H$ and $j_r \Xi_V$.

2.2 Variational sequences

We will be now concerned with some distinguished sheaves of forms on jet spaces [3, 11, 12, 16, 24]. Notice that we will consider sheaves on $J_r \mathbf{Y}$ with respect to the topology generated by open sets of the kind $(\pi_0^r)^{-1}(U)$, with $U \subset \mathbf{Y}$ open in \mathbf{Y} . This is due to the topological triviality of the fibre of $J_r \mathbf{Y} \rightarrow \mathbf{Y}$.

- i. For $r \geq 0$ we consider the standard sheaves $\overset{p}{\Lambda}_r$ of p -forms on $J_r \mathbf{Y}$.
- ii. For $0 \leq s \leq r$ we consider the sheaves $\overset{p}{\mathcal{H}}_{(r,s)}$ and $\overset{p}{\mathcal{H}}_r$ of *horizontal forms*, i.e. of local fibered morphisms over π_s^r and π^r of the type $\alpha : J_r \mathbf{Y} \rightarrow \overset{p}{\wedge} T^* J_s \mathbf{Y}$ and $\beta : J_r \mathbf{Y} \rightarrow \overset{p}{\wedge} T^* \mathbf{X}$, respectively.
- iii. For $0 \leq s < r$ we consider the subsheaf $\overset{p}{\mathcal{C}}_{(r,s)} \subset \overset{p}{\mathcal{H}}_{(r,s)}$ of *contact forms*, i.e. of sections $\alpha \in \overset{p}{\mathcal{H}}_{(r,s)}$ with values into $\overset{p}{\wedge}(\text{im } \vartheta_{s+1}^*)$. We have the distinguished subsheaf $\overset{p}{\mathcal{C}}_r \subset \overset{p}{\mathcal{C}}_{(r+1,r)}$ of local fibered morphisms $\alpha \in \overset{p}{\mathcal{C}}_{(r+1,r)}$ such that $\alpha = \overset{p}{\wedge} \vartheta_{r+1}^* \circ \tilde{\alpha}$, where $\tilde{\alpha}$ is a section of the fibration $J_{r+1} \mathbf{Y} \times_{J_r \mathbf{Y}} \overset{p}{\wedge} V^* J_r \mathbf{Y} \rightarrow J_{r+1} \mathbf{Y}$ which projects down onto $J_r \mathbf{Y}$.

According to [24], the fibered splitting (1) yields the sheaf splitting $\overset{p}{\mathcal{H}}_{(r+1,r)} = \bigoplus_{t=0}^p \overset{p-t}{\mathcal{C}}_{(r+1,r)} \wedge \overset{t}{\mathcal{H}}_{r+1}$, which restricts to the inclusion $\overset{p}{\Lambda}_r \subset \bigoplus_{t=0}^p \overset{p-t}{\mathcal{C}}_r \wedge \overset{t}{\mathcal{H}}_{r+1}^h$, where $\overset{p}{\mathcal{H}}_{r+1}^h := h(\overset{p}{\Lambda}_r)$ for $0 < p \leq n$ and h is defined to be the restriction to $\overset{p}{\Lambda}_r$ of the projection of the above splitting onto the non-trivial summand with the highest value of t . We also define the map $v := \text{id} - h$.

We recall now the theory of variational sequences on finite order jet spaces, as it was developed by Krupka in [11].

By an abuse of notation, let us denote by $d \ker h$ the sheaf generated by the presheaf $d \ker h$ (see [26]). We set $\overset{*}{\Theta}_r := \ker h + d \ker h$.

In [11] it is proved that the following diagram is commutative and that its rows and

columns are exact:

$$\begin{array}{cccccccccccccccc}
 & & 0 & & 0 & & 0 & & 0 & & & & 0 & & 0 & & & & \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & & & \\
 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \overset{1}{\Theta}_r & \xrightarrow{d} & \overset{2}{\Theta}_r & \xrightarrow{d} & \dots & \xrightarrow{d} & \overset{I}{\Theta}_r & \xrightarrow{d} & 0 & \longrightarrow & \dots & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & & & \\
 0 & \longrightarrow & \mathbb{R} & \longrightarrow & \overset{0}{\Lambda}_r & \xrightarrow{d} & \overset{1}{\Lambda}_r & \xrightarrow{d} & \overset{2}{\Lambda}_r & \xrightarrow{d} & \dots & \xrightarrow{d} & \overset{I}{\Lambda}_r & \xrightarrow{d} & \overset{I+1}{\Lambda}_r & \xrightarrow{d} & \dots & \xrightarrow{d} & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & & & \\
 0 & \longrightarrow & \mathbb{R} & \longrightarrow & \overset{0}{\Lambda}_r & \xrightarrow{\varepsilon_0} & \overset{1}{\Lambda}_r / \overset{1}{\Theta}_r & \xrightarrow{\varepsilon_1} & \overset{2}{\Lambda}_r / \overset{2}{\Theta}_r & \xrightarrow{\varepsilon_2} & \dots & \xrightarrow{\varepsilon_{I-1}} & \overset{I}{\Lambda}_r / \overset{I}{\Theta}_r & \xrightarrow{\varepsilon_I} & \overset{I+1}{\Lambda}_r & \xrightarrow{d} & \dots & \xrightarrow{d} & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & & & \\
 & & 0 & & 0 & & 0 & & 0 & & & & 0 & & 0 & & & &
 \end{array}$$

Definition 2.1 The top row of the above diagram is said to be the *contact sequence*, and the bottom row is said to be the *r*-th order *variational sequence* associated with the fibered manifold $\mathbf{Y} \rightarrow \mathbf{X}$. \square

Notice that, in general, the highest integer I depends on the dimension of the fibers of $J_r \mathbf{Y} \rightarrow \mathbf{X}$.

Remark 2.2 If $0 \leq p \leq n$, then $d \ker h \subset \ker h$, and $\alpha \in \ker h$ if and only if $(j_r \sigma)^* \alpha = 0$ for every section $\sigma : \mathbf{X} \rightarrow \mathbf{Y}$; this partly shows the relation of $\overset{p}{\Theta}_r$ to the Calculus of Variations [3, 6, 10, 19].

2.3 Representation of the variational sequence

Isomorphisms of the quotient sheaves with suitable sheaves of forms can be given by means of decomposition formulae.

The sheaf morphism h yields the isomorphisms [23, 24]:

$$(2) \quad I_p : \overset{p}{\Lambda}_r / \overset{p}{\Theta}_r \rightarrow \overset{p}{\mathcal{H}}_{r+1}^h : [\alpha] \mapsto h(\alpha), \quad 0 \leq p \leq n.$$

Let $\beta \in \overset{1}{\mathcal{C}}_r \wedge \overset{n}{\mathcal{H}}_{r+1}^h$. Then there is a unique pair of sheaf morphisms [3, 6, 9, 10, 21, 22]

$$(3) \quad E_\alpha \in \overset{1}{\mathcal{C}}_{(2r,0)} \wedge \overset{n}{\mathcal{H}}_{2r+1}^h, \quad F_\alpha \in \overset{1}{\mathcal{C}}_{(2r,r)} \wedge \overset{n}{\mathcal{H}}_{2r+1}^h,$$

such that $(\pi_{r+1}^{2r+1})^* \alpha = E_\alpha - F_\alpha$ and F_α is locally of the form $F_\alpha = d_H p_\alpha$ with $p_\alpha \in \overset{1}{\mathcal{C}}_{(2r-1,r-1)} \wedge \overset{n-1}{\mathcal{H}}_{2r}$.

The above formula yields the isomorphism

$$(4) \quad I_{n+1} : \overset{n+1}{\Lambda}_r / \overset{n+1}{\Theta}_r \rightarrow \overset{n+1}{\mathcal{V}}_r : [\alpha] \mapsto E_{h(\alpha)}.$$

See [24] for the expression of \mathcal{V}_r^{n+1} .

Let $\eta \in \mathcal{V}_r^{n+1}$. It has been proved [24] that there exists $\tilde{H}_{d\eta} \in \mathcal{C}_{(4r+1,2r+1)}^1 \otimes \mathcal{C}_{(4r+1,0)}^1 \wedge \mathcal{H}_{4r+1}^n$ such that, for all vertical vector fields $\Sigma : \mathbf{Y} \rightarrow V\mathbf{Y}$, we have $E_{\widehat{d\eta}} = \tilde{H}_{d\eta}(j_{2r+1}\Sigma)$, where $\widehat{d\eta} := j_{2r+1}\Sigma \lrcorner d\eta$. Hence there is a unique pair of sheaf morphisms

$$(5) \quad H_{d\eta} \in \mathcal{C}_{(4r+1,2r+1)}^1 \wedge \mathcal{C}_{(4r+1,0)}^1 \wedge \mathcal{H}_{4r+1}^n, \quad G_{d\eta} \in \mathcal{C}_{(4r+1,2r+1)}^2 \wedge \mathcal{H}_{4r+1}^n,$$

such that $(\pi_{2r+1}^{4r+1})^*(d\eta) = H_{d\eta} - G_{d\eta}$ and $H_{d\eta} = \frac{1}{2}A(\tilde{H}_{d\eta})$, where A is the skew-symmetrisation map. Moreover, $G_{d\eta}$ is locally of the type $G_{d\eta} = d_H q_{d\eta}$, where $q_{d\eta} \in \mathcal{C}_{4r}^2 \wedge \mathcal{H}_{4r}^{n-1}$. We recall that $\tilde{H}_{d\eta} = 0$ if and only if $H_{d\eta} = 0$. Remark that, in general, the sections p_α and $q_{d\eta}$ are not uniquely characterized (see, e.g. [3, 9, 24]).

Moreover, we have the isomorphism $I_{n+2} : \mathcal{E}_{n+1}(\mathcal{V}_r^{n+1}) \rightarrow \mathcal{V}_r^{n+2}$, where \mathcal{V}_r^{n+2} is the image of the injective morphism $\mathcal{E}_{n+1}(\Lambda_r^{n+1}/\Theta_r^{n+1}) \rightarrow \mathcal{V}_r^{n+2} : [d\alpha] \mapsto H_{dE_{h(\alpha)}}$.

By means of the above isomorphisms I_p , $0 \leq p \leq n+2$, we obtain a representation of the ‘short’ variational sequence [24], namely

$$0 \longrightarrow \mathbb{R} \longrightarrow \mathcal{V}_r^0 \xrightarrow{\varepsilon_0} \mathcal{V}_r^1 \xrightarrow{\varepsilon_1} \dots \xrightarrow{\varepsilon_n} \mathcal{V}_r^{n+1} \xrightarrow{\varepsilon_{n+1}} \mathcal{V}_r^{n+2} \xrightarrow{\varepsilon_{n+2}} 0.$$

Notice that the morphisms \mathcal{E}_p can be read through the above isomorphisms. In particular, $\mathcal{E}_p(h(\alpha)) = h(d\alpha)$ if $0 \leq p \leq n-1$.

We can interpret the above sequence in terms of the Calculus of Variations (see [3, 6, 9, 16, 24]).

A section $\lambda \in \mathcal{V}_r^n$ is just a Lagrangian of order $(r+1)$ of the standard literature; $\mathcal{E}_n(\lambda) \in \mathcal{V}_r^{n+1}$ is the standard higher order Euler–Lagrange morphism associated with λ . Due to the exactness of the above sequence, if λ is variationally trivial (i.e., $\mathcal{E}_n(\lambda) = 0$ holds), then (locally) $\lambda = d_H \epsilon$, with $\epsilon \in \mathcal{V}_r^{n-1}$.

Let $\alpha \in \Lambda_r^{n+1}$, i.e. $h(\alpha) \in \mathcal{C}_r^1 \wedge \mathcal{H}_{r+1}^n$. We call $E_{h(\alpha)} \in \mathcal{V}_r^{n+1}$ an *Euler–Lagrange type morphism*; it is defined on $J_{2r+1}\mathbf{Y}$. We say $p_{h(\alpha)}$ is a *momentum* associated with $E_{h(\alpha)}$.

Let $\eta \in \mathcal{V}_r^{n+1}$. We call $\mathcal{E}_{n+1}(\eta) = H_{d\eta}$ the *Helmholtz morphism* associated with η ; it is defined on $J_{4r+1}\mathbf{Y}$. Again, the exactness of the sequence implies that, if η is locally variational (i.e., $\mathcal{E}_{n+1}(\eta) = H_{d\eta} = 0 = \tilde{H}_{d\eta}$ holds), then (locally) $\eta = \mathcal{E}_n(\lambda)$ with $\lambda \in \mathcal{V}_r^n$.

Remark 2.3 Let $s \leq r$. Then the inclusions $\Lambda_s^p \subset \Lambda_r^p$ and $\Theta_s^p \subset \Theta_r^p$ yield the injective sheaf morphisms (see [11]) $\chi_s^r : \left(\Lambda_s^p / \Theta_s^p \right) \rightarrow \left(\Lambda_r^p / \Theta_r^p \right) : [\alpha] \mapsto [\pi_s^{r*} \alpha]$, hence the inclusions

$$(6) \quad \kappa_s^r : \mathcal{V}_s^p \rightarrow \mathcal{V}_r^p$$

for $s \leq r$. □

3 Variational Lie derivative

In this section we give a representation of the Lie derivative operator in the variational sequence. We consider a projectable vector field $(\Xi, \bar{\Xi})$ on \mathbf{Y} and take into account the Lie derivative with respect to the prolongation $j_r \Xi$ of Ξ . In fact, such a prolonged vector field preserves the fiberings π_r^s, π^r ; hence it preserves the splitting (1). So, we shall prove that the operator $L_{j_r \Xi}$ passes to the quotient, yielding an operator on the quotient sheaves of the sequence. Moreover, we give the expression of this operator on the representations of the finite order variational sequence.

Representation of Lie derivative

Let $L_{j_r \Xi}$ be the Lie derivative operator with respect to the r -th prolongation $j_r \Xi$ of a projectable vector field $(\Xi, \bar{\Xi})$ on \mathbf{Y} .

As a consequence of earlier results due to Krupka [10], we show that the operator $L_{j_r \Xi}$ defines an operator $\bar{L}_{j_r \Xi}$ on the quotient spaces $\bar{\Lambda}_r / \bar{\Theta}_r$. First we prove a lemma.

Lemma 3.1 *We have*

$$\bar{L}_{j_r \Xi} : \bar{\Lambda}_r / \bar{\Theta}_r \rightarrow \bar{\Lambda}_r / \bar{\Theta}_r : [\alpha] \mapsto \bar{L}_{j_r \Xi}([\alpha]) = [L_{j_r \Xi} \alpha] .$$

PROOF. We have to prove that $L_{j_r \Xi}(\bar{\Theta}_r) \subset \bar{\Theta}_r$. We recall that $\bar{\Theta}_r = \ker h + d \ker h$ and that $\alpha \in d \ker h$ if and only if, locally, $\alpha = d\beta$, where $\beta \in \ker h$. Moreover, it is evident that $L_{j_r \Xi}(d \ker h) = d(L_{j_r \Xi} \ker h)$. So, we have only to prove that $L_{j_r \Xi}(\ker h) \subset \ker h$. To see this, let us consider the inclusion

$$\ker h \subset \bigoplus_{t=0}^q \mathcal{C}_r^{p-t} \wedge \mathcal{H}_{r+1}^t,$$

where $q = p - 1$ if $0 < p \leq n$, and $q = p - n - 1$ if $p > n$.

The above inclusion, together with the standard result [10, 16] $L_{j_r \Xi}(\bar{\mathcal{C}}_r) \subset \bar{\mathcal{C}}_r$, shows that $L_{j_r \Xi}(\ker h)$ has no components in \mathcal{H}_{r+1}^h if $0 \leq p \leq n$, or in $\mathcal{C}_r^{p-n} \wedge \mathcal{H}_{r+1}^n$ if $p > n$.

QED

This allows us to prove

Proposition 3.2 *Let $[\alpha] \in \bar{\Lambda}_r / \bar{\Theta}_r$. The isomorphisms I_p enable us to represent the map $\bar{L}_{j_r \Xi}$ on $\bar{\mathcal{V}}_r$ as follows:*

$$(7) \quad \tilde{\mathcal{L}}_{j_r \Xi} : \bar{\mathcal{V}}_r \rightarrow \bar{\mathcal{V}}_r : I_p([\alpha]) \mapsto \tilde{\mathcal{L}}_{j_r \Xi}(I_p([\alpha])) := I_p(\bar{L}_{j_r \Xi}[\alpha]) = I_p([L_{j_r \Xi} \alpha]) .$$

If we set $h(L_{j_r \Xi} \alpha) = \hat{\alpha}$, then we have

$$\begin{aligned} I_p([\alpha]) &= h(\alpha) \mapsto \hat{\alpha} \quad \text{if } 0 \leq p \leq n, \\ I_p([\alpha]) &= E_{h(\alpha)} \mapsto E_{\hat{\alpha}} \quad \text{if } p = n + 1, \\ I_p([d\alpha]) &= H_{dE_{h(\alpha)}} \mapsto H_{dE_{\hat{\alpha}}} \quad \text{if } p = n + 2. \end{aligned}$$

Definition 3.3 Let $(\Xi, \bar{\Xi})$ be a projectable vector field. We say that the map $\tilde{\mathcal{L}}_{j_r \Xi}$ defined in the above theorem is the *variational Lie derivative*. \square

Noether's theorems

Variational Lie derivatives allow us to calculate infinitesimal symmetries of forms in the variational sequence. In particular, we are interested in symmetries of Lagrangians and Euler–Lagrange morphisms (up to horizontal forms or divergencies). However, the representation in the above theorem is not very useful in concrete cases; here, we provide formulae for calculating symmetries of such morphisms which generalize Noether's theorems.

The inclusions κ_r^s give rise to new representations of $\bar{L}_{j_r \Xi}$ on \mathcal{V}_s^p . We will distinguish between the following cases:

- i. $0 \leq p \leq n - 1$.
- ii. $p = n$, which is of great importance for the Calculus of Variations. In fact, the representation yields the classical Noether's theorem.
- iii. $p = n + 1$, where the representation yields a formula for the symmetries of an Euler–Lagrange type morphism.

Let again $L_{j_r \Xi}$ be the Lie derivative operator with respect to the r -th prolongation $j_r \Xi$ of a projectable vector field $(\Xi, \bar{\Xi})$ on \mathbf{Y} . We start with a technical lemma.

Lemma 3.4 *Let $r \leq s$. We have*

$$\kappa_r^s \circ \tilde{\mathcal{L}}_{j_r \Xi} = \tilde{\mathcal{L}}_{j_s \Xi} \circ \kappa_r^s = I_p \circ \bar{L}_{j_r \Xi} \circ \chi_r^s \circ I_p^{-1} .$$

PROOF. In fact we have $\tilde{\mathcal{L}}_{j_r \Xi} = I_p \circ \bar{L}_{j_r \Xi} \circ I_p^{-1}$, $\kappa_r^s = I_p \circ \chi_r^s \circ I_p^{-1}$. \square

Then the following two results hold true (for $p \leq n$).

Proposition 3.5 *Let $0 \leq p \leq n - 1$ and $I_p([\alpha]) = h(\alpha) \in \mathcal{V}_r^p$. Then we have*

$$\kappa_r^{r+3} \circ \tilde{\mathcal{L}}_{j_r \Xi}(h(\alpha)) = d_H(\bar{\Xi} \lrcorner h(\alpha)) + \bar{\Xi} \lrcorner d_H h(\alpha) + j_{r+2} \Xi_V \lrcorner d_V h(\alpha) .$$

PROOF. The proof can be easily performed by means of the splitting (1), which yields $j_r \Xi \circ \pi_r^{r+1} = j_r \Xi_H + j_r \Xi_V$, $(\pi_r^{r+1})^* \alpha = h(\alpha) + v(\alpha)$ and $(\pi_r^{r+1})^* d = d_H + d_V$. We have

$$\begin{aligned} \kappa_r^{r+3} \circ \tilde{\mathcal{L}}_{j_r \Xi}(h(\alpha)) &= h(L_{j_r \Xi} \alpha) \\ &= h((d_H + d_V)(j_{r+1} \Xi_H + j_{r+1} \Xi_V) \lrcorner h(\alpha)) \\ &\quad + h((j_{r+2} \Xi_H + j_{r+2} \Xi_V) \lrcorner (d_H + d_V)h(\alpha)) \\ &= d_H(j_{r+1} \Xi_H \lrcorner h(\alpha)) + j_{r+2} \Xi_H \lrcorner d_H h(\alpha) + j_{r+2} \Xi_V \lrcorner d_V h(\alpha) . \end{aligned} \quad \square$$

Theorem 3.6 (*First Noether's theorem*) Let $p = n$ and $I_n([\alpha]) = h(\alpha) \in \mathcal{V}_r^n$. Then we have (locally)

$$\kappa_r^{2r+1} \circ \tilde{\mathcal{L}}_{j_r \Xi}(h(\alpha)) = \Xi_V \lrcorner \mathcal{E}(h(\alpha)) + d_H(j_r \Xi_V \lrcorner p_{d_V h(\alpha)} + \bar{\Xi} \lrcorner h(\alpha)) .$$

PROOF. We make use of the same techniques used to prove the above proposition, together with the decomposition (3). We have

$$\begin{aligned} \kappa_r^{2r+1} \circ \tilde{\mathcal{L}}_{j_r \Xi}(h(\alpha)) &= h(L_{j_{r+1} \Xi} h(\alpha)) \\ &= d_H(j_{r+1} \Xi_H \lrcorner h(\alpha)) + h(j_{r+2} \Xi_V \lrcorner d_V h(\alpha)) \\ &= d_H(\bar{\Xi} \lrcorner h(\alpha)) + h(j_{2r+1} \Xi_V \lrcorner (E_{d_V h(\alpha)} + F_{d_V h(\alpha)})) . \end{aligned}$$

Since $F_{d_V h(\alpha)} = d_H p_{d_V h(\alpha)}$ at least locally, because of (3) we obtain (locally)

$$\kappa_r^{2r+1} \circ \tilde{\mathcal{L}}_{j_r \Xi}(h(\alpha)) = \Xi_V \lrcorner \mathcal{E}(h(\alpha)) + d_H(j_r \Xi_V \lrcorner p_{d_V h(\alpha)} + \bar{\Xi} \lrcorner h(\alpha)) . \quad \boxed{QED}$$

Since we are also interested in the case $p = n + 1$, we first prove another technical Lemma.

Lemma 3.7 Let $I_p([\alpha]) \in \mathcal{V}_r^p$ with $p > n$. Then we have (globally)

$$\kappa_r^{r+3} \circ \tilde{\mathcal{L}}_{j_r \Xi} I_p([\alpha]) = [d_H(\bar{\Xi} \lrcorner h(\alpha)) + d_V(j_{r+1} \Xi_V \lrcorner h(\alpha)) + j_{r+2} \Xi_V \lrcorner d_V h(\alpha)] .$$

PROOF. We can easily prove the lemma by making use of the decomposition (5) and of the isomorphisms I_p . We find

$$\begin{aligned} \kappa_r^{r+3} \circ \tilde{\mathcal{L}}_{j_r \Xi} I_p([\alpha]) &= [h(L_{j_r \Xi} \alpha)] \\ &= [h((d_X + d_V)(j_{r+1} \Xi_H + j_{r+1} \Xi_V) \lrcorner h(\alpha)) \\ &\quad + h((j_{r+2} \Xi_H + j_{r+2} \Xi_V) \lrcorner (d_H + d_V)h(\alpha))] \\ &= [d_H(\bar{\Xi} \lrcorner h(\alpha)) + d_V(j_{r+1} \Xi_V \lrcorner h(\alpha)) + j_{r+2} \Xi_V \lrcorner d_V h(\alpha)] , \end{aligned}$$

where we have used $d_H(j_{r+1} \Xi_H \lrcorner h(\alpha)) = d_H(\bar{\Xi} \lrcorner h(\alpha))$. \boxed{QED}

Remark 3.8 As for the case $p = n + 1$, we remark that an Euler–Lagrange type morphism can be given in two ways:

1. as $[\alpha] \in \Lambda_r^{n+1} / \Theta_r^{n+1}$, which yields the morphism $E_{h(\alpha)}$;
2. as $\eta \in \mathcal{V}_r^{n+1}$.

In principle, the two ways are equivalent. But in practical calculations it is very hard to find an $\alpha \in \Lambda_r^{n+1}$ such that $\eta = E_{h(\alpha)}$ holds. \square

Theorem 3.9 (*Second Noether's theorem*) Let $p = n + 1$ and $\alpha \in \Lambda_r^{n+1}$. Then we have

$$\kappa_r^{4r+1} \circ \tilde{\mathcal{L}}_{j_r \Xi} I_{n+1}([\alpha]) = \mathcal{E}(j_r \Xi_V \lrcorner h(\alpha)) + \tilde{H}_{d_{E_{h(\alpha)}}}(j_{2r+1} \Xi_V) .$$

PROOF. By the above lemma we have

$$\begin{aligned} \kappa_r^{4r+1} \circ \tilde{\mathcal{L}}_{j_r \Xi} I_{n+1}([\alpha]) &= I_{n+1}([\mathcal{E}(j_r \Xi_V \lrcorner h(\alpha)) + j_{r+2} \Xi_V \lrcorner d_V h(\alpha)]) \\ &= [\mathcal{E}(j_r \Xi_V \lrcorner h(\alpha)) + j_{2r+1} \Xi_V \lrcorner d_V E_{h(\alpha)}] \\ &= \mathcal{E}(j_r \Xi_V \lrcorner h(\alpha)) + \tilde{H}_{dE_{h(\alpha)}}(j_{2r+1} \Xi_V) . \end{aligned}$$

□ QED

Remark 3.10 Let $p = n + 1$ and $I_{n+1}([\eta]) = \eta \in \mathcal{V}_r^{n+1} \subset \mathcal{V}_{2r+1}^{n+1}$. Then we have

$$\kappa_{2r+1}^{4r+1} \circ \tilde{\mathcal{L}}_{j_r \Xi} \eta = \mathcal{E}(\Xi_V \lrcorner \eta) + \tilde{H}_{d\eta}(j_{2r+1} \Xi_V) .$$

Noether's theorems and variational sequences

Here we summarize the above results on symmetries in the finite order variational sequence.

Theorem 3.11 Let $(\Xi, \bar{\Xi})$ be a projectable vector field on \mathbf{Y} and $r \leq s$. We have the commutative diagram

$$\begin{array}{ccc} \mathcal{V}_r^p & \xrightarrow{\mathcal{L}_{j_r \Xi}} & \mathcal{V}_s^p \\ & \searrow \mathcal{L}_{j_r \bar{\Xi}} & \nearrow \kappa_r^s \\ & \mathcal{V}_r^p & \end{array}$$

where $\mathcal{L}_{j_r \Xi}$ is defined as follows:

1. if $0 \leq p \leq n - 1$ and $\mu \in \mathcal{V}_r^p$, then $s = r + 3$ and

$$\mathcal{L}_{j_r \Xi}(\mu) = d_H(\bar{\Xi} \lrcorner \mu) + \bar{\Xi} \lrcorner d_H \mu + j_{r+2} \Xi_V \lrcorner d_V \mu ;$$

2. if $p = n$ and $\lambda \in \mathcal{V}_r^n$, then $s = 2r + 1$ and

$$\mathcal{L}_{j_r \Xi}(\lambda) = \Xi_V \lrcorner \mathcal{E}(\lambda) + d_H(j_r \Xi_V \lrcorner p_{d_V} \lambda + \bar{\Xi} \lrcorner \lambda) ;$$

3. if $p = n + 1$ and $\alpha \in \Lambda_r^{n+1}$, then $s = 4r + 1$ and

$$\mathcal{L}_{j_r \Xi}(E_{h(\alpha)}) = \mathcal{E}(j_r \Xi_V \lrcorner h(\alpha)) + \tilde{H}_{dE_{h(\alpha)}}(j_{2r+1} \Xi_V) .$$

4 Relations with the Calculus of Variations

In this section we analyse some consequences of Noether's theorems. In particular, we find the relationship between abstract Noether's theorems and some standard concepts of the Calculus of Variations. The above theorems play an important role in mathematical and physical applications concerning "variationally relevant" symmetries of Lagrangians and Euler–Lagrange morphisms.

Definition 4.1 Let $(\Xi, \bar{\Xi})$ be a projectable vector field on \mathbf{Y} . Let $\lambda \in \mathcal{V}_r^n$ be a Lagrangian and $\eta \in \mathcal{V}_r^{n+1}$ an Euler–Lagrange morphism. Then Ξ is called a *symmetry* of λ (of η) if $\mathcal{L}_{j_{r+1}\Xi} \lambda = 0$ (respectively, $\mathcal{L}_{j_{2r+1}\Xi} \eta = 0$). \square

Remark 4.2 Due to $\mathcal{E}\mathcal{L}_{j_r\Xi} = \mathcal{L}_{j_r\Xi}\mathcal{E}$, a symmetry of a Lagrangian λ is also a symmetry of its Euler–Lagrange morphism E_λ (but the converse is not true, see e.g. [10, 13, 18]). \square

Let now $\eta \in \mathcal{V}_r^{n+1}$ be an Euler–Lagrange morphism and let $\sigma : \mathbf{X} \rightarrow \mathbf{Y}$ be a section. We recall that σ is said to be *critical* if $\eta \circ j_{2r+1}\sigma = 0$.

Remark 4.3 Let $\lambda \in \mathcal{V}_r^n$ be a Lagrangian and $(\Xi, \bar{\Xi})$ a symmetry of λ . Then, by theorem 3.6, i.e. the first Noether's theorem, we have

$$0 = \Xi_V \lrcorner \mathcal{E}(\lambda) + d_H(j_r \Xi_V \lrcorner p_{d_V \lambda} + \bar{\Xi} \lrcorner \lambda).$$

Suppose that the section $\sigma : \mathbf{X} \rightarrow \mathbf{Y}$ fulfils $(j_{2r+1}\sigma)^*(\Xi_V \lrcorner \mathcal{E}(\lambda)) = 0$. This condition holds for all critical sections, but it is not equivalent to the evolution equations since, in principle, it can hold also on some non-critical sections. Then the $(n-1)$ -form $\epsilon = j_r \Xi_V \lrcorner p_{d_V \lambda} + \bar{\Xi} \lrcorner \lambda$ of \mathcal{V}_{2r}^{n-1} fulfils the equation

$$(8) \quad d((j_{2r}\sigma)^*(j_r \Xi_V \lrcorner p_{d_V \lambda} + \bar{\Xi} \lrcorner \lambda)) = 0.$$

If σ is a critical section the above equation (8) admits a physical interpretation as a conservation law along critical sections for the density associated with ϵ . \square

Definition 4.4 Let $\lambda \in \mathcal{V}_r^n$ be a Lagrangian and Ξ a symmetry of λ . Then a sheaf morphism of the type

$$\epsilon = (j_r \Xi_V \lrcorner p_{d_V \lambda} + \bar{\Xi} \lrcorner \lambda) \in \mathcal{V}_{2r}^{n-1}$$

is said to be a *conserved current*. \square

Remark 4.5 In general, a conserved current is not uniquely defined. In fact, it depends on the choice of $p_{d_V \lambda}$. Moreover, we could add to the conserved current any form $\beta \in \mathcal{V}_{2r}^{n-1}$ which is variationally closed, i.e. such that $\mathcal{E}_{n-1}(\beta) = 0$ holds. Furthermore, β is locally of the type $\beta = d_H \gamma$, where $\gamma \in \mathcal{V}_{2r+1}^{n-2}$. \square

In the sequel we give some preliminary examples for the first and second order Lagrangians making use of the coordinate expressions given in [24].

Remark 4.6 Let $r = 1$ and $\lambda \in \overset{n}{\mathcal{H}}_1 \subset \overset{n}{\mathcal{V}}_1$. Then we have the coordinate expression $\lambda = L\omega$, where $L \in \overset{0}{\Lambda}_1$. Hence we have the following expression for the conserved current:

$$\epsilon = ((\Xi^i - y_\gamma^i \Xi^\gamma) \partial_i^\mu L + \Xi^\mu L) \omega_\mu. \quad \square$$

Remark 4.7 Let $r = 2$ and $\lambda \in \overset{n}{\mathcal{H}}_2 \subset \overset{n}{\mathcal{V}}_2$. Then we have the coordinate expression $\lambda = L\omega$, where $L \in \overset{0}{\Lambda}_2$. Hence we have the following expression for the conserved current:

$$\epsilon = ((\Xi^i - y_\gamma^i \Xi^\gamma) (\partial_i^\mu L - D_\nu \partial_i^{\mu\nu} L) + D_\nu (\Xi^i - y_\gamma^i \Xi^\gamma) \partial_i^{\mu\nu} L + \Xi^\mu L) \omega_\mu. \quad \square$$

Similar considerations can be made for the Euler–Lagrange morphisms.

Remark 4.8 Let $\eta \in \overset{n+1}{\mathcal{V}}_r$ and let Ξ be a symmetry of η . Then, by Remark 3.10, we have

$$0 = \mathcal{E}(\Xi_V \lrcorner \eta) + \tilde{H}_{d\eta}(j_{2r+1} \Xi_V).$$

Suppose that η is locally variational, i.e. $H_{d\eta} = \tilde{H}_{d\eta} = 0$; then we have

$$(9) \quad \mathcal{E}(\Xi_V \lrcorner \eta) = 0.$$

This implies that $\Xi_V \lrcorner \eta$ is variationally trivial. $\Xi_V \lrcorner \eta$ is locally of the type $\Xi_V \lrcorner \eta = d_H \beta$, where $\beta \in \overset{n-1}{\mathcal{V}}_{r+1}$.

Suppose that the section $\sigma : \mathbf{X} \rightarrow \mathbf{Y}$ fulfils $(j_{2r+1} \sigma)^*(\Xi_V \lrcorner \eta) = 0$. Then we have $d((j_{2r} \sigma)^* \beta) = 0$ so that, as in the case of Lagrangians, if σ is a critical section, then β is conserved along σ . \square

Definition 4.9 Let $\eta \in \overset{n+1}{\mathcal{V}}_r$ be an Euler–Lagrange morphism and Ξ a symmetry of η . Then a sheaf morphism of the type $\beta \in \overset{n-1}{\mathcal{V}}_{r+1}$ fulfilling the conditions of the above remark is called a *conserved current*. \square

Remark 4.10 As in the case of Lagrangians, a conserved current for an Euler–Lagrange morphism is not uniquely defined. In fact, we could add to $\Xi_V \lrcorner \eta$ any variationally trivial Lagrangian, obtaining different conserved currents. Moreover, such conserved currents are defined up to variationally trivial $(n-1)$ -forms. \square

Remark 4.11 In general, it is difficult to find a conserved current β of the above type. A possible way to find β is the following one. If η is locally variational, it is possible to find a (local) Lagrangian λ such that $\mathcal{E}(\lambda) = \eta$. Then the conserved current is found with the same procedure as for Lagrangians. We notice that the problem of finding the Lagrangian (inverse problem) is rather difficult (see [11, 12, 17, 24]). \square

5 An example of physical relevance

In this section we will consider as an example of application the case of the gravitational field interacting with an external “matter” field [4]. We shall show how the formalism developed here enables us to obtain in a very straightforward way well known results about conserved quantities in General Relativity in presence of “matter”.

Let \mathbf{Y} be a *natural* bundle (see *e.g.* [7] for a review of naturality). The geometrical framework concerning *natural symmetries* will be developed in detail in a forthcoming paper [5]. Here it is enough to recall that the Lie derivative of a section of \mathbf{Y} (see *e.g.* [4, 7]) can be locally written as

$$\mathcal{L}_\xi y^i = y^i_\gamma \xi^\gamma - \mathcal{Z}_\gamma^{i\alpha} \xi^\gamma,$$

where α is a multi-index of length $|\alpha| = k$ (the order of the natural bundle) and $\mathcal{Z}_\lambda^{i\alpha} \in C^\infty(\mathbf{Y})$. Moreover, in this case $-\Xi_V$ is just the Lie derivative of sections of the bundle \mathbf{Y} .

The following coordinate expressions hold:

$$d_V \lambda = (d_V \lambda)_i^\alpha \vartheta_\alpha^i \wedge \omega, \quad E_{d_V \lambda} = \mathcal{E}(\lambda)_i \vartheta^i \wedge \omega, \quad p_{d_V \lambda} = p(\lambda)_i^{\alpha\mu} \vartheta_\alpha^i \wedge \omega_\mu.$$

It is known (see *e.g.* [9]) that the decomposition formula (3) when applied to $d_V \lambda$ gives

$$(10) \quad (d_V \lambda)_i^\alpha = p(\lambda)_i^{\beta\mu} \quad \beta + \mu = \alpha, |\alpha| = r,$$

$$(11) \quad (d_V \lambda)_i^\alpha = p(\lambda)_i^{\beta\mu} + D_\nu p(\lambda)_i^{\alpha\nu} \quad \beta + \mu = \alpha, |\alpha| = r - 1,$$

$$(12) \quad (d_V \lambda)_i^\alpha = \mathcal{E}(\lambda)_i^\alpha + D_\nu p(\lambda)_i^{\alpha\nu} \quad |\alpha| = 0.$$

Furthermore, $\mathcal{E}(\lambda)_i = \sum_{|\alpha| \leq r} (-1)^{|\alpha|} D_\alpha (d_V \lambda)_i^\alpha$.

Let us now take $\mathbf{Y} = \underset{\mathbf{X}}{\text{Lor}}(\mathbf{X}) \times \mathbf{F}$, where $\text{Lor}(\mathbf{X})$ is the bundle of Lorentzian metrics over the space-time \mathbf{X} and \mathbf{F} is a natural bundle of “matter” fields [4]. Let us consider the *natural* Lagrangian λ defined on the bundle $J_2 \mathbf{Y}$:

$$(13) \quad \lambda = \lambda(g^{\mu\nu}, \gamma_{\mu\nu}^\alpha, R_{\mu\nu}, \varphi^\Lambda, \varphi_\mu^\Lambda)$$

$$(14) \quad = \lambda_H(g^{\mu\nu}, R_{\mu\nu}) + \lambda_M(g^{\mu\nu}, \gamma_{\mu\nu}^\alpha, \varphi^\Lambda, \varphi_\mu^\Lambda)$$

$$(15) \quad = \lambda_H(j_2 g) + \lambda_M(j_1 g; j_1 \varphi),$$

where $\lambda_H = -\frac{1}{2\kappa} \sqrt{g} g^{\alpha\beta} R_{\alpha\beta}$ is the Hilbert Lagrangian, $R_{\alpha\beta}$ is the (formal) Ricci tensor of the metric g given by $R_{\alpha\beta} := R_{\alpha\mu\beta}^\mu = D_\mu \gamma_{\alpha\beta}^\mu - D_\beta \gamma_{\alpha\mu}^\mu + \gamma_{\nu\mu}^\mu \gamma_{\alpha\beta}^\nu - \gamma_{\nu\beta}^\mu \gamma_{\alpha\mu}^\nu$, with $\gamma_{\nu\beta}^\mu = \frac{1}{2} g^{\mu\alpha} (D_\nu g_{\beta\alpha} - D_\alpha g_{\nu\beta} + D_\beta g_{\alpha\nu})$ the (formal) Levi-Civita connection of g , $\sqrt{g} = \sqrt{|\det(g^{\mu\nu})|}$, κ is a constant and $\lambda_M(j_1 g; j_1 \varphi)$ is the “matter” Lagrangian describing the dynamics of “matter” fields φ interacting with the gravitational field. Then the index i stands for the set of indices $\{\mu\nu, \Lambda\}$ with $\mu, \nu = 1, \dots, \dim \mathbf{X}$ and $\Lambda = 1, \dots, \dim \mathbf{F}$.

By a direct computational method, we can compare Equations (10)–(12) with results already obtained *e.g.* in [4]. In fact, let us make the substitutions

$$\begin{aligned} (d_V \lambda)_\Lambda &= f_\Lambda + p_\Theta^\mu \gamma_{\beta\mu}^\alpha \partial_\Lambda \mathcal{Z}_\alpha^{\Theta\beta}, \\ (d_V \lambda)_\Lambda^\mu &= p(\lambda)_\Lambda^\mu = p_\Lambda^\mu, \\ (d_V \lambda)_{\mu\beta}^\alpha &= \frac{1}{2}(p_{\Lambda\mu} \mathcal{Z}_\beta^{\Lambda\alpha} + p_{\Lambda\beta} \mathcal{Z}_\mu^{\Lambda\alpha}) \equiv A_{\beta\mu}^\alpha, \\ (d_V \lambda)_{\alpha\beta} &= -\frac{1}{2\kappa} \sqrt{g} g_{\alpha\beta} = p(\lambda)_{\alpha\beta}, \\ (d_V \lambda)_{\mu\nu} &= \frac{1}{2} t_{\mu\nu} - \frac{1}{2\kappa} \sqrt{g} (R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu}), \end{aligned}$$

where $R = g^{\alpha\beta} R_{\alpha\beta}$ is the Ricci scalar curvature and we set $p_i^{0\nu} := p_i^\nu$ for the convenience of notation.

It is easy to verify that Equation (8) with the above mentioned substitutions gives

$$\begin{aligned} \epsilon^\sigma &= -\frac{1}{2\kappa} \sqrt{g} g^{\alpha\beta} \mathcal{L}_\xi u_{\alpha\beta}^\sigma + T_{\mu\nu}^\sigma \mathcal{L}_\xi g^{\mu\nu} + p_\Lambda^\sigma \mathcal{L}_\xi \varphi^\Lambda - \xi^\sigma \lambda \\ &\quad + \nabla_\mu \left[\frac{1}{2\kappa} \sqrt{g} (\nabla^\sigma \xi^\mu - \nabla^\mu \xi^\sigma) \right] \\ &\quad + E_\nu^\sigma \xi^\nu + E_\nu^{\sigma\mu} \nabla_\mu \xi^\nu - T_\nu^\sigma \xi^\nu + T_\nu^{\sigma\mu} \nabla_\mu \xi^\nu + (d_V \lambda)_\nu^\sigma \xi^\nu. \end{aligned}$$

Here $u_{\alpha\beta}^\sigma = \gamma_{\alpha\beta}^\sigma - \frac{1}{2} [\delta_\alpha^\sigma \gamma_{\nu\beta}^\nu + \delta_\beta^\sigma \gamma_{\nu\alpha}^\nu]$, $\nabla_\nu \varphi^\Lambda = D_\nu \varphi^\Lambda + \mathcal{Z}_\alpha^{\Lambda\beta}(\varphi) \gamma_{\beta\nu}^\alpha$ and $T_{\mu\nu}^\sigma$ is defined in terms of p_Λ^σ by means of (11) and (12) in the following way:

$$\begin{aligned} 2T_{\mu\nu}^\sigma &= -2[p_\Lambda^\sigma \mathcal{Z}_{(\mu\nu)}^\Lambda + p_{\Lambda(\mu} \mathcal{Z}_{\nu)}^{\Lambda\sigma}] - [p_{\Lambda\nu} \mathcal{Z}_\mu^{\Lambda\sigma} + p_{\Lambda\mu} \mathcal{Z}_\nu^{\Lambda\sigma}] \\ &= -2A_{(\mu\nu)}^\sigma - A_{\nu\mu}^\sigma = p(\lambda)_{\mu\nu}^\sigma. \end{aligned}$$

The tensor density $T_{\mu\nu}^\sigma$ is related by (12) to the so-called *Hilbert stress tensor* density $T_{\mu\nu} = t_{\mu\nu} - \nabla_\sigma T_{\mu\nu}^\sigma$ [4]. Furthermore, the tensorial coefficients E_ν^σ and $E_\nu^{\sigma\mu}$ are defined by

$$E_\nu^\sigma = (d_V \lambda_M)_\Lambda^\sigma \nabla_\nu \varphi^\Lambda - \delta_\nu^\sigma \lambda_M, \quad E_\nu^{\sigma\mu} = -(d_V \lambda_M)_\Lambda^\sigma \mathcal{Z}_\nu^{\Lambda\mu}(\varphi),$$

and are called *energy–momentum tensors* of the theory.

Remark 5.1 This approach shows clearly that the density $T_{\mu\nu}^\sigma$ expresses the coupling of the matter field with the derivatives of the metric, because it defines the relationship between the momentum $p(\lambda)_{\mu\nu}^\sigma$ “associated” to the gravitational Lagrangian and the momentum $p(\lambda)_\Lambda^\sigma$ “associated” to the matter Lagrangian. \square

6 Conclusions

The introduction of the variational Lie derivative enables us to collect a wide range of concepts and coordinate formulae. Namely, we obtain a natural framework from the

idea that Lie derivatives with respect to prolonged vector fields preserve the contact subsequence.

The representation of the variational sequence yields explicit formulae for the variational Lie derivative. Moreover, we obtain expressions for the conserved currents in a straightforward way. Furthermore, our approach enables us to know which is the degree of arbitrariness [19] when we deal with conserved currents (see Remarks 4.5 and 4.10).

Making use of this natural framework, in a forthcoming paper [5] we shall give a geometrical interpretation of the superpotentials in natural field theories (see [4] and references quoted therein). An extension to gauge–natural field theories will be considered in [2].

References

- [1] I. M. ANDERSON, T. DUCHAMP: On the existence of global variational principles, *Amer. Math. J.* **102** (1980) 781–868.
- [2] L. FATIBENE, M. FRANCAVIGLIA, M. PALESE: Conservation laws and variational sequences in gauge–natural theories, to appear in *Math. Proc. Cambridge Phil. Soc.* **130** (2) (2001) xx–xx.
- [3] M. FERRARIS: Fibered connections and global Poincaré–Cartan forms in higher–order calculus of variations, in: *Proc. Diff. Geom. and its Appl.* (Nové Město na Moravě, 1983); D. Krupka ed., J.E. Purkyně University (Brno, 1984) 61–91.
- [4] M. FERRARIS, M. FRANCAVIGLIA: The Lagrangian approach to conserved quantities in general relativity, *Mechanics, Analysis and Geometry: 200 Years after Lagrange*; M. Francaviglia ed., Elsevier Science Publishers B. V. (Amsterdam 1991), 451–488.
- [5] M. FRANCAVIGLIA, M. PALESE, R. VITOLO: Superpotentials in variational sequences, *Proc. VII Conf. Diff. Geom. and Appl., Satellite Conf. of ICM in Berlin* (Brno 1998); I. Kolář *et al.* eds.; Masaryk University in Brno (Czech Republic) 1999, 469–480.
- [6] P. L. GARCIA, J. MUÑOZ: On the geometrical structure of higher order variational calculus, *Proc. IUTAM–ISIMM Symp. on Modern Developments in Anal. Mech.*, (Torino, 1982); S. Benenti, M. Francaviglia, A. Lichnerowicz eds., Tecnoprint (Bologna, 1983) 127–147.
- [7] I. KOLÁŘ, P.W. MICHOR, J. SLOVÁK: *Natural Operations in Differential Geometry*, (Springer–Verlag, N.Y., 1993).
- [8] I. KOLÁŘ: Lie derivatives and higher order Lagrangians, *Proc. Diff. Geom. and its Appl.* (Nové Město na Moravě, 1980); O. Kowalski ed., Univerzita Karlova (Praha, 1981) 117–123.
- [9] I. KOLÁŘ: A geometrical version of the higher order hamilton formalism in fibred manifolds, *J. Geom. Phys.* **1** (1984) (2) 127–137.
- [10] D. KRUPKA: Some geometric aspects of variational problems in fibred manifolds, *Folia Fac. Sci. Nat. UJEP Brunensis* **14**, J. E. Purkyně Univ. (Brno, 1973) 1–65.
- [11] D. KRUPKA: Variational sequences on finite order jet spaces, *Proc. Diff. Geom. and its Appl.* (Brno, 1989); J. Janyška, D. Krupka eds., World Scientific (Singapore, 1990) 236–254.
- [12] D. KRUPKA: Topics in the calculus of variations: finite order variational sequences, *Proc. Diff. Geom. and its Appl.* (Opava, 1993) 473–495.
- [13] D. KRUPKA, A. TRAUTMAN: General invariance of Lagrangian structures, *Bull. Acad. Polon. Sci., Math. Astr. Phys.* **22** (1974) (2) 207–211.

- [14] L. MANGIAROTTI, M. MODUGNO: Fibered spaces, jet spaces and connections for field theories, in *Proc. Int. Meet. on Geom. and Phys.*, Pitagora Editrice (Bologna, 1983) 135–165.
- [15] J. NOVOTNÝ: Modern methods of differential geometry and the conservation laws problem; *Folia Fac. Sci. Nat. UJEP Brunensis (Physica)* **19** (1974) 1–55.
- [16] D. J. SAUNDERS: *The Geometry of Jet Bundles*, Cambridge Univ. Press (Cambridge, 1989).
- [17] F. TAKENS: A global version of the inverse problem of the calculus of variations, *J. Diff. Geom.* **14** (1979) 543–562.
- [18] A. TRAUTMAN: Noether equations and conservation laws, *Comm. Math. Phys.* **6** (1967) 248–261.
- [19] A. TRAUTMAN: A metaphysical remark on variational principles, *Acta Phys. Pol. B* **XX** (1996) 1–9.
- [20] W. M. TULCZYJEW: The Lagrange complex, *Bull. Soc. Math. France* **105** (1977) 419–431.
- [21] A. M. VINOGRADOV: On the algebro–geometric foundations of Lagrangian field theory, *Soviet Math. Dokl.* **18** (1977) 1200–1204.
- [22] A. M. VINOGRADOV: A spectral sequence associated with a non–linear differential equation, and algebro–geometric foundations of Lagrangian field theory with constraints, *Soviet Math. Dokl.* **19** (1978) 144–148.
- [23] R. VITOLO: On different geometric formulations of Lagrangian formalism, *Diff. Geom. and its Appl.* **10** (1999) 225–255.
- [24] R. VITOLO: Finite order Lagrangian bicomplexes, *Math. Proc. Cambridge Phil. Soc.* **125** (1) (1998) 321–333.
- [25] R. VITOLO: A new infinite order formulation of variational sequences, *Arch. Math. Univ. Brunensis* **34** (1998) 483–504.
- [26] R. O. WELLS: *Differential Analysis on Complex Manifolds*, *GTM*, n. **65**, Springer–Verlag (Berlin, 1980).

Authors' addresses:

Mauro Francaviglia, Marcella Palese
 Department of Mathematics, University of Turin
 Via C. Alberto 10, 10123 Turin, Italy
 E-mail: francaviglia@dm.unito.it, palese@dm.unito.it

Raffaele Vitolo
 Department of Mathematics “E. De Giorgi”, University of Lecce
 Via per Arnesano, 73100 Lecce, Italy
 E-mail: vitolo@ilenic.unile.it